



# Tau Neutrino Searches at Super-Kamiokande and Hyper-Kamiokande

Roger Wendell  
Kyoto University  
NuTau 2021  
2021.09.29




SK




HK

# Super-Kamiokande:



- 22.5 kton fiducial volume
- Optically separated into
  - Inner Detector 11,146 20" PMTs
  - Outer Detector 1885 8" PMTs
- No net electric or magnetic fields
- Excellent PID between showering (e-like) and non-showering ( $\mu$ -like)
  - $< 1\%$  MIS ID at 1 GeV
- Multipurpose physics

Concentration vs Capture Eff. on Gd



**Four Run Periods:**  
SK-I (1996-2001) SK-II (2003-2005)  
SK-III (2005-2008) **SK-IV (2008-2018)**  
**SK-V (2019-2020)**

**Upgrade Complete Now operating as (SK-Gd) !!**

# Hyper-Kamiokande:

- **186.5 kton** fiducial volume (258 kton total)
- Optically separated into
  - Inner Detector
    - **20,000** 50cm High QE B&L PMTs
    - + O(1) k mPMT Modules
  - Outer Detector 1885 8" PMTs
- Pure water neutron tagging > 40%
- Now **under construction** – Data taking in 2027




50 cm HQE  
Box&Line PMT

SK QE  $\times 2$   
SK  $\sigma_\tau \times 2$   
SK Pres. Tol  $\times 2$




Multi-PMT



Current Hyper-K analysis uses the same methodology as Super-K,  
In 10 years of data expect  
→ A factor of **~6 times** the current SK data set  
→ No improvements assumed in systematic errors or detector performance...yet

# Atmospheric Neutrino Flux:

PHYSICAL REVIEW D 94, 052001 (2016)





No  $\nu_\tau$  in the flux at SK/HK energies



# Atmospheric Neutrino Flux: $\nu_\mu$ Disappearance

PHYSICAL REVIEW D 94, 052001 (2016)





$$P_{\nu_\mu \rightarrow \nu_\tau} \simeq \cos^2 \theta_{13} \sin^2(2\theta_{23}) \sin^2 \left( 1.27 \Delta m_{32}^2 \frac{L}{E} \right)$$

$\text{Log}_{10}(E_\nu/\text{GeV})$

Super-Kamiokande

Well known that this best explained by  $\nu_\mu \rightarrow \nu_\tau$  oscillations

# Tau Expectation:




- Assuming current values of oscillation parameters expect roughly (CC)

$$\Gamma_{\tau} \sim 1/\text{yr}/\text{kton}$$



- Current SK data set spans 1996-2016 representing 328 kton-yrs
- Assumed **HK** data set 1.86 **M**ton-yrs

Impact of  $v_\tau$

# Hyper-Kamiokande : Mass Hierarchy Prospects



Multi-GeV e-like  $\nu_e$





- Tau neutrinos are expected to appear in the same kinematic region as the mass hierarchy signal
- Uncertainty in the tau cross section (currently using  $\sim 25\%$ ) has a large impact on expected sensitivity
- $\rightarrow$  Secondary impact on q23 octant sensitivity

# Future : Other Physics with Atmospheric Neutrinos

*J. Gehrlein, TAUP 2021*

*Hyper-K Design Report*




- Core-crossing atmospheric neutrinos can probe electron density of the core via its impact on their oscillations
  - General test of earth matter effects – not living on a vacuum, probably
- High-statistics observations can be used to **constrain the electron content of the Earth**
  - Important in geophysics for understanding the origin of the Earth's magnetic field
- Tau appearance contribution to (non-)unitarity studies



# Searching for $\nu_\tau$

3 Flavor  $P(\nu_\mu \rightarrow \nu_\tau)$




CC  $\nu_\tau$

Super-Kamiokande IV

Run 999999 Sub 2 Event 7  
16-04-13:05:43:18  
Inner: 8104 hits, 30188 pe  
Outer: 3 hits, 2 pe  
Trigger: 0x07  
D\_walls: 1130.7 cm  
Evis: 3.3 GeV

Charge (pe)


- >26.7
- 23.3-26.7
- 20.2-23.3
- 17.3-20.2
- 14.7-17.3
- 12.2-14.7
- 10.0-12.2
- 8.0-10.0
- 6.2- 8.0
- 4.7- 6.2
- 3.3- 4.7
- 2.2- 3.3
- 1.3- 2.2
- 0.7- 1.3
- 0.2- 0.7
- < 0.2




- Complicated event topologies with many overlapping charged particles

# Search for Tau Neutrinos at SK

## Hadronic Decay



## Leptonic Decay




Visible Final State identical to that of  
a primary  $\nu_l$


- A 10 GeV  $\tau$  will travel about 0.5 mm in tank before decaying
- Leptonic decay
  - Swamped by charged current backgrounds from other atmospheric neutrinos,  $\nu_l$
- Hadronic decay
  - Expect more isotropic distribution of final state particles
  - Due to production threshold many interactions are DIS, so additional forward-going hadrons

# Search for Tau Neutrinos at SK

CCQE + Hadronic




CC DIS + Hadronic



- A 10 GeV  $\tau$  will travel about 0.5 mm in tank before decaying
- Leptonic decay
  - Swamped by charged current backgrounds from other atmospheric neutrinos,  $\nu_l$
- Hadronic decay
  - Expect more isotropic distribution of final state particles
  - Due to production threshold many interactions are DIS, so additional forward-going hadrons


# Search for Tau Neutrinos at SK: Selection



- Pre-selection
  - Fully-Contained
  - Fiducial Volume (22.5kton)
  - **Visible Energy > 1330.0 MeV**
- Seven Neural Network Variables
  - Log10 (Visible Energy)
  - Particle ID Likelihood
  - Number of Decay Electrons
  - Distance to farthest electron
  - Event sphericity
  - Number of Ring Candidates
  - Fraction of momentum in leading ring

- Select  $\nu_\tau$  above production threshold

# Search for Tau Neutrinos at SK: Selection




- Dependent upon reconstruction to find most energetic ring
- Many tau decays produce showering particles (via e or  $\pi^0$ )
  - CC  $\nu_\mu$  likely to have non-showering



# Search for Tau Neutrinos at SK: Selection

\* Data points are downward-going atm.  $\nu$



$$S^{\alpha\beta} = \frac{\sum_i p_i^\alpha p_i^\beta}{\sum_i p_i^2}$$

$$S = \frac{3}{2} (\lambda_2 + \lambda_3)$$


- Pre-selection
  - Fully-Contained
  - Fiducial Volume (22.5kton)
  - Visible Energy > 1330.0 MeV
- Seven Neural Network Variables
  - Log10 (Visible Energy)
  - Particle ID Likelihood
  - Number of Decay Electrons
  - Distance to farthest electron
  - **Event sphericity**
  - Number of Ring Candidates
  - Fraction of momentum in leading ring

- Calculated based on distribution of charge seen in event COM
- $S = 0$  : linear event ,  $S = 1$  isotropic charge
- $\rightarrow$  Decay of heavy  $\tau$  produces more isotropic charge

# Search for Tau Neutrinos at SK: Selection


## Super-Kamiokande IV

Run 999999 Sub 2 Event 1195  
18-02-07:23:43:29  
Inner: 8524 hits, 43967 pe  
Outer: 5 hits, 5 pe  
Trigger: 0x07  
D\_wall: 430.2 cm  
Evis: 3.9 GeV




## Charge (pe)

- >26.7
- 23.3-26.7
- 20.2-23.3
- 17.3-20.2
- 14.7-17.3
- 12.2-14.7
- 10.0-12.2
- 8.0-10.0
- 6.2- 8.0
- 4.7- 6.2
- 3.3- 4.7
- 2.2- 3.3
- 1.3- 2.2
- 0.7- 1.3
- 0.2- 0.7
- < 0.2



Recon struction  
result shown as  
cyan circles




- Pre-selection
  - Fully-Contained
  - Fiducial Volume (22.5kton)
  - Visible Energy > 1330.0 MeV
- Seven Neural Network Variables
  - Log10 (Visible Energy)
  - Particle ID Likelihood
  - Number of Decay Electrons
  - Distance to farthest electron
  - Event sphericity
  - **Number of Ring Candidates**
  - Fraction of momentum in leading ring

- Difficult to fully reconstruct all particles, so count ring pieces
- Typically  $\tau$  decay produces more pions above C threshold

# Search for Tau Neutrinos at SK: Selection



## Super-Kamiokande IV

Run 999999 Sub 2 Event 1195  
18-02-07:23:43:29  
Inner: 8524 hits, 43967 pe  
Outer: 5 hits, 5 pe  
Trigger: 0x07  
D\_wall: 430.2 cm  
Evis: 3.9 GeV



## Charge (pe)

- >26.7
- 23.3-26.7
- 20.2-23.3
- 17.3-20.2
- 14.7-17.3
- 12.2-14.7
- 10.0-12.2
- 8.0-10.0
- 6.2-8.0
- 4.7-6.2
- 3.3-4.7
- 2.2-3.3
- 1.3-2.2
- 0.7-1.3
- 0.2-0.7



## Pre-selection

- Fully-Contained
- Fiducial Volume (22.5kton)
- Visible Energy > 1330.0 MeV


## Seven Neural Network Variables

- Log10 (Visible Energy)
- Particle ID Likelihood
- Number of Decay Electrons
- Distance to farthest electron
- Event sphericity
- **Number of Ring Candidates**
- Fraction of momentum in leading ring


Count ring pieces

- Typically  $\tau$  decay produces more pions above C threshold

# Search for Tau Neutrinos at SK: Selection



$$= \frac{\sum_{\theta_i < 48^\circ} q_i}{\sum q_i}$$




- Pre-selection
  - Fully-Contained
  - Fiducial Volume (22.5kton)
  - Visible Energy > 1330.0 MeV
- Seven Neural Network Variables
  - Log10 (Visible Energy)
  - Particle ID Likelihood
  - Number of Decay Electrons
  - Distance to farthest electron
  - Event sphericity
  - Number of Ring Candidates
  - Fraction of momentum in leading ring

- Calculate charge with 48 degrees of highest momentum particle
- Expect more even energy distribution in  $\nu_\tau$  events → small fractions

# Search for Tau Neutrinos at SK: Performance

PHYS. REV. D 98, 052006 (2018)





NN > 0.5	Background	Signal ( $\nu_\tau$ )
Efficiency	28%	76%
Purity	95.3%	4.7% ←
Rate [Mton· year] <sup>-1</sup>	8467	422



# Search for Tau Neutrinos at SK: Performance


PHYS. REV. D 98, 052006 (2018)



NN > 0.5	Background	Signal ( $\nu_\tau$ )	Decay mode	Branching ratio (%)	Tau-like fraction (%)
Efficiency	28%	76%	$e^- \bar{\nu}_e \nu_\tau$	17.83	$67.3 \pm 2.2$
Purity	95.3%	4.7%	$\mu^- \bar{\nu}_\mu \nu_\tau$	17.41	$42.6 \pm 2.6$
Rate [Mton·year] <sup>-1</sup>	8467	422	$\pi^- \nu_\tau$	10.83	$84.7 \pm 3.8$
			$\pi^- \pi^0 \nu_\tau$	25.52	$81.0 \pm 2.1$
			$3\pi \nu_\tau$	18.29	$88.7 \pm 2.5$
			Others	10.12	$90.5 \pm 3.4$

# Search for Tau Neutrinos at SK :

PHYS. REV. D **98**, 052006 (2018)




$$\text{Data} = \text{PDF}_{\text{BG}} + \alpha \times \text{PDF}_{\text{tau}} + \sum \epsilon_i \times \text{PDF}_i$$

$$\alpha = 1.47 \pm 0.32 \quad (\text{stat+syst})$$

4.6 $\sigma$  rejection of no  $\tau$  appearance

- Fit 2-dimensional PDFs ( $\cos \theta$ , Neural Network ), while simultaneously varying systematic error templates
  - No cut to separate tau-like and non-tau-like
- Uses 328 kton-yr exposure (SK-I ~ SK-IV)

# Systematic Errors in Search for Tau Neutrinos



■ Flux-averaged cross section:


$$\sigma_{measured} = (1.47 \pm 0.32) \times \langle \sigma_{theory} \rangle$$

■ Consistent with model at about  $1\sigma$

$$= (0.94 \pm 0.20) \times 10^{-38} \text{ cm}^2$$

Stat+Syst.

# Systematic Errors in Search for Tau Neutrinos




- Dominant systematics are cross section related but flux errors also play a role
- Total systematic uncertainty is  $\sim 8\%$  impact on normalization
- Present analysis is dominated by statistical uncertainty  $\sim 20\%$

*In decreasing order of impact*

Systematic error	$\sigma$ (%)	
NC/CC ratio	20%	20
DIS $q^2$ dependence for low W	15%	10
Meson exchange current		10
$1\pi$ axial coupling		10
DIS $q^2$ dependence for high W		10
Coherent $\pi$ cross section		100
Flux normalization ( $E_\nu > 1\text{GeV}$ )	11%	15
$1\pi$ background scale factor		10
$1\pi$ axial form factor		10
CCQE cross section		10
Single pion $\pi^0/\pi^\pm$ ratio		40
$\bar{\nu}_\mu/\nu_\mu$ ratio ( $E_\nu > 10\text{ GeV}$ )	5.6%	15
$\nu/\bar{\nu}$ ratio ( $E_\nu > 10\text{ GeV}$ )	5.3%	5
DIS cross section ( $E_\nu < 10\text{ GeV}$ )		10
FC multi-GeV normalization		5
$\bar{\nu}_e/\nu_e$ ratio ( $E_\nu > 10\text{GeV}$ )	4.6%	8
$K/\pi$ ratio	4.5%	10
Single meson cross section		20
Single-pion $\bar{\nu}/\nu$ ratio		10
Horizontal/vertical ratio	3%	1
CCQE $\nu/\bar{\nu}$ ratio		10
DIS cross section		5
Matter effect		6.8
Neutrino path length		10

\* Numbers in red are maximum impact on analysis bins, red boxes are flux-related errors





# Expanding Analysis to Hyper-Kamiokande



Hyper-Kamiokande-like  
 $\nu_\tau$  MC

20% B&L Coverage  
+5k mPMT equiv.

Charge (pe)




\* Same event as above, viewed in HK-like display

- Simple scaling of analysis above to 5.6 Mton-yr
  - ... or 30 years of Hyper-K
- Corresponds to  $\sim 8\%$  uncertainty on normalization  $\rightarrow$  limit of current assumptions



# Prospects and Ideas

- Neutron Information
  - Tau events: more hadrons → more neutrons
  - Another handle on isotropy?
  - *Unknown and potentially large systematics*
- Improved reconstruction algorithm
  - SK Fiducial volume 22.5 → 27.0<sup>+</sup> kton
  - Better ring identification
  - Better allotment of charge among rings
  - “New” algorithm is current default for HK
- Improved multi-variate methods
  - Graduate from TMVA...
- Hyper-K specific tools are under development
  - Take advantage of improve QE PMTs
  - Complementary information from mPMTs
  - Machine Learning-based reconstruction



True Number of Rings	New Reco.			SK Standard		
	1R	2R	≥ 3R	1R	2R	≥ 3R
True 1R	95.0%	4.64%	0.41%	95.9%	3.85%	0.29%
True 2R	27.8%	66.7%	5.56%	42.5%	52.8%	4.63%
True ≥3R	7.04%	25.5%	67.5%	20.2%	33.0%	46.8%

*Reconstruction efficiency*

# Summary and Conclusions

- Tau neutrinos appearing in the SK atmospheric neutrino data at  $4.6\sigma$ 
  - Currently statistics limited measurement
- At Hyper-K level exposures sensitivity to reach 8% error on tau normalization
  - Assumes SK analysis → currently being ported to HK specific tools
- Prospects for improved reconstruction, additional NN variables, are expected to improve the S/N for short-term and can be extended in principle to long-term
- Techniques presented here can equally be applied to HK beam neutrinos from J-PARC
  - Stay tuned....

# Super-Kamiokande collaboration



Kamioka Observatory, ICRR, Univ. of Tokyo, Japan  
 RCCN, ICRR, Univ. of Tokyo, Japan  
 University Autonoma Madrid, Spain  
 University of British Columbia, Canada  
 Boston University, USA  
 University of California, Irvine, USA  
 California State University, USA  
 Chonnam National University, Korea  
 Duke University, USA  
 Fukuoka Institute of Technology, Japan  
 Gifu University, Japan  
 GIST, Korea

University of Hawaii, USA  
 Imperial College London, UK  
 NFN Bari, Italy  
 INFN Napoli, Italy  
 INFN Padova, Italy  
 INFN Roma, Italy  
 Kavli IPMU, The Univ. of Tokyo, Japan  
 KEK, Japan  
 Kobe University, Japan  
 Kyoto University, Japan  
 University of Liverpool, UK  
 LLR, Ecole polytechnique, France  
 Miyagi University of Education, Japan

ISEE, Nagoya University, Japan  
 NCBJ, Poland  
 Okayama University, Japan  
 Osaka University, Japan  
 University of Oxford, UK  
 Queen Mary University of London, UK  
 Seoul National University, Korea  
 University of Sheffield, UK  
 Shizuoka University of Welfare, Japan  
 Sungkyunkwan University, Korea  
 Stony Brook University, USA  
 Tokai University, Japan  
 The University of Tokyo, Japan

Tokyo Institute of Technology, Japan  
 Tokyo University of Science, Japan  
 University of Toronto, Canada  
 TRIUMF, Canada  
 Tsinghua University, Korea  
 The University of Winnipeg, Canada  
 Yokohama National University, Japan

178 collaborators  
 from 45 institutes  
 10 countries





京大学  
UNIVERSITY OF TOKYO

ハイパーカミオカンデ 着工記念式典

Hyper-Kamiokande Groundbreaking Ceremony



宇宙線研

Institute for Cosmic Ray Research, The University of Tokyo